

2-2017

The Effects of Closed Kinetic Chain and Endurance Exercises on Reducing Pain in a Child with Ehlers-Danlos Syndrome: A Case Report

Linda Duong

University of St. Augustine for Health Sciences

Megan Flores

University of St. Augustine for Health Sciences, mflores@usa.edu

Follow this and additional works at: <https://soar.usa.edu/pt>



Part of the [Physical Therapy Commons](#)

Recommended Citation

Duong, Linda and Flores, Megan, "The Effects of Closed Kinetic Chain and Endurance Exercises on Reducing Pain in a Child with Ehlers-Danlos Syndrome: A Case Report" (2017). *Physical Therapy Collection*. 16.

<https://soar.usa.edu/pt/16>

This Conference Proceeding is brought to you for free and open access by the Faculty and Staff Research at SOAR @ USA. It has been accepted for inclusion in Physical Therapy Collection by an authorized administrator of SOAR @ USA. For more information, please contact soar@usa.edu.

The Effects of Closed Kinetic Chain and Endurance Exercises on Reducing Pain in a Child with Ehlers-Danlos Syndrome: A Case Report

Linda Duong, PT, DPT^{1,2}; Megan Flores, PT, MPT, PCS¹

¹University of St. Augustine for Health Sciences, Doctor of Physical Therapy Program

Background and Purpose

- Children with Ehlers-Danlos Syndrome hypermobility type (EDS) are prone to developing long-term musculoskeletal pain and fatigue from joint laxity.¹
- Physical therapy with emphasis on weight-bearing exercises can improve stability and proprioception in individuals with hypermobile joints.^{2,3}

The purpose of this case report was to describe the effectiveness of closed kinetic chain (CKC) and endurance exercises on reducing pain in a child with EDS.

Case Description

Subject: 9-year-old girl with EDS (hypermobility type)
PT Diagnosis: bilateral posterior tibialis tendonitis

International Classification of Functioning model⁴:



Methods

- Six-week physical therapy program focused on dynamic stability and improving joint control
- 45-60 minute sessions, twice a week
- Use of a Lower Body Positive Pressure Treadmill (LBPPT) to address cardiovascular endurance⁵
- Child and family education on proper body mechanics and joint protection



www.performbetter.com



www.alterg.com

Exercises Utilized During Treatment Sessions:

Targeted Exercises (Weeks Performed)	Exercise Type	Purpose
Ankle Strengthening (Weeks 1-2)	Toe-heel raises, towel scrunches, marble pick-ups, foot intrinsic series	Strengthening of ankle dorsiflexors, plantarflexors, foot intrinsics
Flexibility (Weeks 1-6)	Hamstring stretches, incline board stretches	Improve lower extremity flexibility in tight muscles
Lower Extremity Strengthening (Weeks 1-6)	Bridges on swiss ball, mini wall squats, terminal knee extensions and sidesteps with theraband, leg press	Incorporate weight-bearing exercises to engage multiple joints and muscle co-contraction
Balance Training (Weeks 2-6)	Tiltboard, circleboard, single leg balance, tandem walking (even and uneven surfaces)	Improve balance and proprioception, static/dynamic stabilization, neuromuscular re-education
Cardiovascular Training (Weeks 3-6)	Recumbent bike, LBPPT	Improve endurance, improve high-impact weight-bearing tolerance

LBPPT = Lower Body Positive Pressure Treadmill

Results

Evaluation:

- Lower Extremity Functional Scale (LEFS) score of 42/80
- Reported pain of 7/10 on Faces Pain Scale
- Child wore ankle stabilizing orthotics at all times
- Knee hyperextension posture in standing
- Unable to participate in sports and running

Discharge:

- Increased LEFS score to 53/80
- Decreased pain to 2/10 on Faces Pain Scale
- Reduced need for ankle stabilizing orthotics (only worn during sports)
- Neutral knee extension posture in standing
- Child returned to sports and running

Discussion

- Interventions which utilize CKC exercises can help children with EDS achieve more functional outcomes.⁶
- The use of a LBPPT can allow children with joint pain to improve their cardiovascular endurance by progressing weight-bearing tolerance over time.^{5,7}
- The child in this case report demonstrated improved strength, proprioception, and balance through participation in physical therapy.
- The child and parents reported satisfaction with overall mobility and safe return to recreational activities.

Conclusion

- The findings from this case report support the use of strengthening and endurance exercises to improve joint stability and prevent further injury in children with EDS.
- Future studies should investigate additional treatment strategies towards promoting functional outcomes in this population.

References

1. Cma C, Action S. Identification and management of joint hypermobility. 2014:1-22.
2. Simpson MR. Benign joint hypermobility syndrome: evaluation, diagnosis, and management. *J Am Osteopat Asso.* 2006;106(9):531-536.
3. Nikolajsen H, Larsen PK, Simonsen EB, et al. Gait pattern in 9-11-year-old children with generalized joint hypermobility compared with controls; a cross-sectional study. *BMC Musculoskelet Disord.* 2013;14:341.
4. APTA. Guide to Physical Therapist Practice 3.0. Alexandria, VA. 2014. <http://guidetoptpractice.apta.org/>. Accessed May 23, 2016.
5. El-refay BH, Faiad NT. Body Composition Response to Lower Body Positive Pressure Training in Obese Children. 2014;8(8):468-473.
6. Kemp S, Roberts I, Gamble C, et al. A randomized comparative trial of generalized vs targeted physiotherapy in the management of childhood hypermobility. *Rheumatology.* 2010;49(2):315-325.
7. Damiano D, DeJong S. A systematic review of the effectiveness of treadmill training and body weight support in pediatric rehabilitation. *J Neurol Phys Ther.* 2009;33(1):27-44.