

1-1-2022

## Lower Extremity and Trunk Electromyographic Muscle Activity During Performance of the Y-Balance Test on Stable and Unstable Surfaces.

Navpreet Kaur

*University of St. Augustine for Health Sciences, nkaur@usa.edu*

Kunal Bhanot

*University of St. Augustine for Health Sciences, kbhanot@usa.edu*

Germaine Ferreira

*University of St. Augustine for Health Sciences, gferreira@usa.edu*

Follow this and additional works at: <https://soar.usa.edu/pt>



Part of the [Physical Therapy Commons](#), and the [Sports Medicine Commons](#)

---

### Recommended Citation

Kaur N, Bhanot K, Ferreira G. Lower Extremity and Trunk Electromyographic Muscle Activity During Performance of the Y-Balance Test on Stable and Unstable Surfaces.. *International Journal of Sports Physical Therapy*. 2022; 17(3):483-492. doi: <https://doi.org/10.26603/001c.32593>

This Article is brought to you for free and open access by the Faculty and Staff Research at SOAR @ USA. It has been accepted for inclusion in Physical Therapy Collection by an authorized administrator of SOAR @ USA. For more information, please contact [soar@usa.edu](mailto:soar@usa.edu), [erobinson@usa.edu](mailto:erobinson@usa.edu).

Original Research

# Lower Extremity and Trunk Electromyographic Muscle Activity During Performance of the Y-Balance Test on Stable and Unstable Surfaces

Navpreet Kaur<sup>1 a</sup>, Kunal Bhanot<sup>2</sup>, Germaine Ferreira<sup>1</sup><sup>1</sup> Physical Therapy, University of St. Augustine for Health Sciences, Austin, TX, <sup>2</sup> Physical Therapy, Carlow University, Pittsburgh, PA

Keywords: y balance test, star excursion balance test, electromyography, dynamic balance

<https://doi.org/10.26603/001c.32593>

---

## International Journal of Sports Physical Therapy

Vol. 17, Issue 3, 2022

---

### Background

The Star Excursion Balance Test (SEBT) has been used as a rehabilitation exercise. To improve its efficacy, efficiency, and method variations, the Y-Balance Test (YBT) with anterior (A), posterolateral (PL), and posteromedial (PM) directions of the SEBT has been recommended. Electromyographic activity has been reported to change when the same task is performed on various surfaces.

### Hypothesis/Purpose

To compare the EMG activity of trunk and LE muscles during the performance of the YBT on stable and unstable surfaces.

### Study Design

Cross-Sectional study.

### Methods

Healthy adults with no history of chronic ankle instability were recruited for the study. Surface electromyography was collected for bilateral (ipsilateral [i] and contralateral [c]) rectus abdominis (RA), external oblique (EOB), erector spinae (ES). While, gluteus maximus (GMAX), gluteus medius (GMED), medial hamstrings (MH), biceps femoris (BF), vastus medialis (VM), rectus femoris (RF), vastus lateralis (VL), anterior tibialis (AT), and medial gastrocnemius (MG) on the stance leg (ipsilateral side), during the performance of the YBT. The unstable surface was introduced using a Thera-Band stability trainer. Differences in electromyography were examined for each reach direction and muscle between the stable and unstable surfaces ( $p \leq 0.05$ ).

### Results

Twenty (10 male, 10 female) subjects participated (age:  $27.5 \pm 4.0$  years, height:  $167 \pm 1.0$  cm, weight:  $66.5 \pm 13.0$  kg, body fat:  $14.1 \pm 6.2\%$ ). Significantly higher muscle activity for the unstable surface ( $p < 0.05$ ) with moderate to large effect sizes were observed for the following muscles in the A direction: GMED, GMAX, VM, RF, and MG; PL direction: iEOB, iES, cES, GMED, BF, VM, RF, and MG; and PM direction iEOB, iES, GMED, BF, VM, and RF. Significantly higher muscle activity for the stable surface ( $p = 0.007$ ) was observed in MH muscle in the A direction. No significant differences ( $p > 0.05$ ) between the stable and

---

<sup>a</sup> **Corresponding author:**

Dr. Navpreet Kaur, PT, PhD  
Associate Professor,  
Doctor of Physical Therapy Program,  
University of St. Augustine for Health Sciences,  
5401 LaCrosse Avenue, Austin, TX, 78739  
Phone Number: 737-202-3338  
Fax Number: 512-394-9764  
Email: [nkaur@usa.edu](mailto:nkaur@usa.edu)

unstable surfaces were observed in iRA, cRA, cEOB, VL, and AT for any of the directions of the YBT.

## Conclusion

An increase in muscle activity was observed during YBT on unstable versus stable surfaces for some muscles.

## Level of Evidence

2B

## INTRODUCTION

The Y-Balance Test (YBT), derived from the Star Excursion Balance Test (SEBT), is an inexpensive and commonly used objective measure to assess lower extremity (LE) dynamic balance, functional symmetry, and stability.<sup>1-4</sup> Due to the possibility of errors that may occur during the performance of the SEBT and the absence of a standardized protocol for the performance of the test, the YBT has been gaining popularity in clinical and research settings.<sup>3-5</sup> The YBT was designed to standardize the SEBT test performance and to improve the reliability of the test. The YBT™ (Functional Movement Screen, Danville, VA) is an instrumented version of the modified SEBT that assesses an individual's performance in the anterior (A), posterolateral (PL), and posteromedial (PM) directions. To perform the YBT, the individual stands on an elevated central plastic platform and pushes a rectangular reach block with the foot along a calibrated plastic tubing in each of the three directions.

The YBT has been documented as a reliable tool to assess balance in athletes.<sup>3,6</sup> Smith et al.<sup>7</sup> found that A reach direction asymmetry on the YBT was associated with an increased risk for injury across various sports. They concluded that the YBT could be a valuable tool when screening athletes across multiple sports. Similar results were found by Butler et al.<sup>8</sup> who suggested that poor performance on the YBT is associated with an increased risk for LE injuries in college football players. Poor performance on the YBT has also been linked to an increased incidence of ankle injuries in collegiate athletes.<sup>9</sup> Increased risk of injuries has also been documented in high school cross-country male runners who demonstrated greater asymmetries while performing the YBT.<sup>1</sup>

It seems reasonable that identifying the risk factors associated with sports-related injuries could help clinicians/trainers/coaches in designing programs/interventions to address those factors and overall decrease the injury rates. LE injuries have been attributed to poor neuromuscular (NM) control.<sup>5,7,10</sup> Neuromuscular control is defined as the detection, perception, and utilization of relevant sensory information to perform specific tasks.<sup>11</sup> To reduce injury risk, training programs should include diverse interventions to improve dynamic balance and control. The SEBT has been utilized as a training tool to improve trunk NM control and dynamic balance assessed by measuring trunk strength and endurance using a pressure biofeedback unit and with single leg stance time with eyes open and closed, respectively.<sup>12,13</sup> Significant improvements in these outcomes were seen in the subjects who were trained using the SEBT. No such studies have been published utilizing the YBT tool kit as a training instrument.

Another way of improving dynamic balance is by introducing unstable surfaces to the training programs.<sup>14,15</sup> Unstable surfaces cause a greater challenge to NM control in comparison to stable surfaces and hence are commonly used by clinicians and trainers in their protocols. While introducing instability to exercises may improve stability, the effect on muscle activation has not been completely described. Surface EMG has been extensively used to measure muscle activity (activation) during various activities. Muscle activation, as measured using surface electromyography (EMG), has been documented to vary significantly depending on the type of surface.<sup>14-16</sup> Adding instability may be effective in increasing the muscle activation in stabilizing muscles.<sup>15</sup>

Rehabilitation professionals commonly use the intensity of exercise, defined as a given percentage of the maximal muscle contraction and measured by the EMG, to design protocols for strength and stability training.<sup>16-18</sup> Previous researchers have documented parameters to determine sufficient EMG activation to achieve various training effects.<sup>17,19,20</sup> Strength gains are expected from exercises that cause EMG activation levels greater than 40% however, activation levels below 40% are still beneficial in improving NM control.<sup>17,19,20</sup> Authors have found EMG activation to be direction-specific when measured in subjects performing the SEBT.<sup>19,21,22</sup> To the authors' knowledge, no studies measuring the EMG activity while performing the YBT have been published. Knowing the muscle activation patterns while performing YBT on various surfaces may assist in utilizing this test as a training tool. Therefore, the purpose of this study was to compare the EMG activity of trunk and LE muscles during the YBT on stable and unstable surfaces.

## METHODS

### PARTICIPANTS

Twenty healthy adults (10 males and 10 females) were recruited via email to the university community to participate in the study. The study was approved by the Institutional Review Board of the University of St. Augustine for Health Sciences, Austin, TX.

### PROTOCOL

The participants signed the informed consent before starting the test protocol. The participants were screened for the inclusion/exclusion criteria ([Table 1](#)).<sup>19</sup> Lange® skinfold calipers (model # 68902, Fitness Mart®, division of Country Technology, Inc. Gays Mills, WI) and three site formula regression equations for men (chest, abdomen, and thigh) and women (triceps, supra-iliac, and abdomen) were used

**Table 1. Inclusion/exclusion criteria of the study.**

Inclusion Criteria	Exclusion Criteria
<ul style="list-style-type: none"> <li>• Age 18-40 years,</li> <li>• Age-related body composition (% body fat) between fair to very lean, as reported in ACSM's Guidelines for Exercise Testing and Prescription <ul style="list-style-type: none"> <li>◦ Men: Age 20-29: 4.2 - 18.6 %body fat; Age 30-39: 7.3 - 21.6 %body fat.</li> <li>◦ Women: Age 20-29: 11.4 - 23.5 %body fat; Age 30-39: 11.0 - 24.8 %body fat</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>• History of chronic ankle instability (CAI) of the stance leg (the leg participants would stand on to kick a ball)</li> <li>• Upper or lower extremity injury within prior 6 months</li> <li>• History of upper extremity surgery within prior 6 months</li> <li>• Any history of neck, back, or lower extremity surgery.</li> <li>• Currently experiencing pain anywhere in the body</li> <li>• Difficulty maintaining single leg stance for 10 seconds on either leg</li> <li>• Visible contra-lateral pelvic drop during single-leg stance</li> <li>• History of head injury or any other disorder affecting their balance.</li> </ul>

to assess body composition.<sup>23</sup> Skinfold measurements were taken by the guidelines provided by the *ACSM's Guidelines for Exercise Testing and Prescription*<sup>23</sup> at the sites mentioned above for both the men and the women. The body composition of the participants was assessed to achieve the most accurate surface EMG (sEMG) signal by reducing the effects of body fat on the sEMG signal by excluding participants with body fat percentage above the "Fair" category by age (Men: Age 20-29: 18.6 %body fat; Age 30-39: 21.6 %body fat. Women: Age 20-29: 23.5 %body fat; Age 30-39: 24.8 %body fat).<sup>24</sup>

The Trigno wireless EMG system (Delsys Inc. Boston, MA, USA) was used to collect all the EMG data. The wireless electrodes used measured 37mm x 26mm x 15 mm and encompassed a hard-wired single differential amplifier and a four-bar (99.9% silver) contact area, with an inter-electrode distance of 10 mm (Delsys Inc. Boston, MA, USA). Surface EMG was collected from the erector spinae (ES), external oblique (EOB), and rectus abdominis (RA) bilaterally (ipsilateral and contralateral side of the stance leg). While, gluteus medius (GMED), gluteus maximus (GMAX), medial hamstrings (MH), biceps femoris (BF), vastus medialis (VM), rectus femoris (RF), vastus lateralis (VL), anterior tibialis (AT), and medial gastrocnemius (MG) on the stance leg during the YBT. The skin was cleaned with a skin prep pad, and the area was shaved if body hair was present. The electrodes were placed according to the procedure described by Cram et al.<sup>25</sup> The participants performed light jumping jacks for 30 seconds for warm-up after the electrodes were placed.<sup>26</sup> For normalization of the EMG data, maximum voluntary isometric contractions (MVIC) were performed for each muscle. The MVIC test positions were consistent with those demonstrated by Kendall<sup>27</sup> and previous research.<sup>17,19</sup> Proper electrode placements were also confirmed by observing the EMG amplitudes during the manual muscle tests before testing the MVIC. The participants were asked to perform primary muscle action of the neighboring muscles to assess for crosstalk. Manual pressure was gradually increased until maximum resistance was applied and then held for five seconds using a metronome. Each muscle test was repeated three times with the thirty-second rest between contractions. For all subjects, MVIC was averaged across the three intermediate seconds for each muscle to calculate the mean of the peak RMS value of the three trials.

Two minutes of rest was provided between the MVIC testing of different muscles.

#### Y-BALANCE PERFORMANCE

A YBT kit™ was placed on the floor. This requires participants to stand on one limb and to perform a reaching task in three directions with the other lower limb while maintaining balance on the stance leg ([Figure 1](#)).<sup>3,4,6,10</sup> The preferred stance leg was defined as the leg participant would stand on to kick a ball because it would simulate unilateral weight-bearing activities of the participants and also to make comparisons with previous EMG studies performed during the SEBT.<sup>19</sup> Participants were asked to place the foot of their stance leg behind the designated line on the central platform on the tool kit. Participants completed the test barefoot. Participants were instructed to keep their arms by the side so their shoulder flexion and abduction did not exceed 45° while performing the YBT. No specific instructions were provided for trunk motion. An unstable surface was introduced by placing a Theraband™ stability trainer on top of the central platform ([Figure 2](#)). The order of the reach directions and surface conditions were randomized using computerized random sequence generator.

Participants were instructed to toe touch on the central platform before beginning to reach, marking the beginning of the single-leg stance phase. The toe touch event was recognized on the EMG data using a toe sensor that showed a signal spike on the computer screen. Participants were instructed to push the rectangular reach block as far as possible along the calibrated plastic tubing in one of the predetermined directions of the YBT with the opposite leg and return to the double-leg stance. They were asked to toe touch on the central platform before putting any weight on the reaching leg marking the end of the single-leg stance phase. A metronome was used at a rate of 30 beats/min (equates to two seconds) to ensure consistent timing and speed of each of the YBT trials where each reaching phase (from initial stance to maximum reach; two seconds) and the recovery phase (from maximum reach to bilateral stance; two seconds) was performed during one beat, a total of four seconds.<sup>19</sup> Verbal cueing was provided to the participants during the YBT to synchronize their toe touch with the metronome beat.

Maximum reach distances were recorded at the touch-down point. The reach distance was normalized by participants' leg length measured from the anterior superior iliac spine (ASIS) to the distal end of the medial malleolus.<sup>28</sup> Participants completed six practice trials with the metronome in each of the three (A, PL, PM) reach directions of the YBT. A five-minute break was provided between the practice trials and the data collection. Fifteen seconds of recovery time was given between test trials to reduce the risk of fatigue. A 60 second recovery time was utilized between reach directions, and the order of the reach directions was randomized. The trial was discarded if the heel of the stance leg lifted off the ground, or if the participant put weight on the reaching leg during maximal reach, lost balance even if the heel remained on the ground, could not return to the starting position, or did not match the metronome speed. If the trial was discarded, additional trials were performed until the participant completed three good trials in each direction.

#### DATA PROCESSING

The data were collected at a sampling frequency of 1926 Hz, common-mode reduction ratio > 80 dB@60 Hz; signal to noise ratio of > 750 nv, and no gain was applied to the signal. All collected signals were subsequently bandpass filtered (between 20 and 450 Hz) with a 2<sup>nd</sup> order filter on the high-pass and a 4<sup>th</sup> order filter on the low-pass, then rectified and finally smoothed by using a root-mean-square (RMS) calculation. RMS was calculated using a default window length of 0.125 s with a 0.0625s window overlap. MVIC was used to calculate the percentage MVIC. The mean RMS value of the EMG signal of each muscle for each direction during the YBT was calculated during the two seconds reaching (eccentric) phase of each YBT trial. The reaching phase was measured from the beginning of the unilateral stance to the maximal reach identified by the toe sensor. The EMG of the reaching phase was calculated to be consistent with the methods described in the prior literature.<sup>19,22</sup> The RMS value of the reaching phase of the three trials was averaged for each muscle to be normalized to its respective MVIC value and represented as a percentage of the MVIC (%MVIC).


$$\%MVIC = \left( \frac{\text{Average RMS value of the three repetitions}}{\text{average peak RMS value}} \right) \times 100$$

#### STATISTICAL ANALYSIS


Paired t-tests were used to determine differences in %MVIC of the three reach directions A (stable vs. unstable), PL (stable vs. unstable), and PM (stable vs. unstable) for each muscle. The level of significance was pre-set at  $p \leq 0.05$ , and SPSS version 23.0 was used for all the statistical analyses. The Cohens d effect size was also calculated for each comparison.<sup>29</sup>

#### RESULTS

Twenty (10 male, 10 female) subjects participated (age:  $27.5 \pm 4.0$  years, height:  $167 \pm 1.0$  cm, weight:  $66.5 \pm 13.0$  kg, body fat:  $14.1 \pm 6.2\%$ ). Significant differences with moder-



**Figure 1. Participant demonstrating the Y-Balance Test in the posterolateral direction on the stable surface.**



**Figure 2. Participant demonstrating the Y-Balance Test in the posterolateral direction on the unstable surface.**

ate to large effect sizes between stable and unstable surfaces (greater EMG values in the unstable condition) were observed for the following muscles in the A direction:

**Table 2. EMG activity of each muscle presented as %MVIC (maximal voluntary isometric contraction).**

Directions Muscles	A			PL			PM		
	Stable (Mean ± SD)	Unstable (Mean ± SD)	Effect Size	Stable (Mean ± SD)	Unstable (Mean ± SD)	Effect Size	Stable (Mean ± SD)	Unstable (Mean ± SD)	Effect Size
iRA	13.1 ± 11.0	14.2 ± 12.4	0.14	7.1 ± 6.2	8.5 ± 9.0	0.43	6.1 ± 4.6	7.2 ± 6.9	0.35
cRA	9.0 ± 6.1	8.9 ± 5.3	0.04	5.1 ± 3.7	5.8 ± 5.0	0.4	5.2 ± 3.8	5.9 ± 4.9	0.43
iEOB	18.6 ± 15.1	22.1 ± 20.5	0.4	<b>12.7 ± 9.3*</b>	<b>20.6 ± 20.6*</b>	<b>0.6</b>	<b>10.8 ± 9.9*</b>	<b>14.0 ± 14.1*</b>	<b>0.6</b>
cEOB	32.4 ± 37.7	40.7 ± 53.5	0.5	25.0 ± 36.8	29.3 ± 35.3	0.4	36.8 ± 44.7	42.0 ± 46.8	0.4
iES	19.7 ± 17.1	22.3 ± 18.1	0.25	<b>63.2 ± 27.5*</b>	<b>69.7 ± 31.6*</b>	<b>0.6</b>	<b>27.4 ± 13.9*</b>	<b>33.1 ± 15.5*</b>	<b>0.5</b>
cES	21.2 ± 17.7	23.9 ± 16.7	0.33	<b>27.6 ± 12.2*</b>	<b>32.4 ± 15.6*</b>	<b>0.6</b>	36.1 ± 14.1	39.0 ± 16.5	0.45
GMED	<b>35.5 ± 25.7*</b>	<b>42.2 ± 23.0*</b>	<b>0.96</b>	<b>23.4 ± 15.1*</b>	<b>27.5 ± 18.0*</b>	<b>0.6</b>	<b>36.5 ± 19.2*</b>	<b>42.7 ± 21.6*</b>	<b>0.9</b>
GMAX	<b>10.5 ± 6.4*</b>	<b>12.5 ± 9.3*</b>	<b>0.5</b>	12.9 ± 7.6	13.0 ± 7.2	0.03	11.8 ± 6.0	12.8 ± 6.6	0.2
MH	<b>30.8 ± 17.5*</b>	<b>22.8 ± 13.3*</b>	<b>0.7</b>	18.1 ± 10.4	19.8 ± 12.0	0.2	19.3 ± 10.3	19.2 ± 11.2	0.02
BF	19.3 ± 10.9	20.5 ± 14.0	0.2	<b>23.6 ± 12.9*</b>	<b>27.7 ± 12.8*</b>	<b>0.6</b>	<b>15.3 ± 8.8*</b>	<b>19.2 ± 13.2*</b>	<b>0.7</b>
VM	<b>97.1 ± 57.8*</b>	<b>127.8 ± 56.7*</b>	<b>1.2</b>	<b>63.2 ± 27.2*</b>	<b>73.8 ± 30.8*</b>	<b>0.9</b>	<b>89.5 ± 39.7*</b>	<b>100.5 ± 46.6*</b>	<b>0.7</b>
RF	<b>32.0 ± 22.8*</b>	<b>55.5 ± 32.0*</b>	<b>1.6</b>	<b>33.2 ± 27.1*</b>	<b>40.8 ± 29.6*</b>	<b>1.0</b>	<b>42.4 ± 32.8*</b>	<b>51.4 ± 30.4*</b>	<b>1.1</b>
VL	<b>88.5 ± 38.6*</b>	<b>116.1 ± 46.6*</b>	<b>1.6</b>	<b>65.9 ± 32.8*</b>	<b>77.4 ± 37.8*</b>	<b>1.2</b>	<b>87.4 ± 44.4*</b>	<b>96.1 ± 48.4*</b>	<b>0.7</b>
AT	41.6 ± 18.5	43.9 ± 12.5	0.2	52.1 ± 17.0	55.3 ± 19.2	0.3	47.5 ± 19.2	46.6 ± 18.3	0.08
MG	<b>38.8 ± 20.5*</b>	<b>48.9 ± 24.3*</b>	<b>0.7</b>	<b>47.9 ± 27.8*</b>	<b>58.3 ± 32.7*</b>	<b>0.9</b>	41.2 ± 27.6	44.8 ± 26.1	0.3

SD, Standard Deviation; A, Anterior; PM, Posteromedial; PL, Posterolateral; iRA, Ipsilateral rectus abdominis; cRA, Contralateral rectus abdominis; iEOB, Ipsilateral external oblique; cEOB, Contralateral external oblique; iES, Ipsilateral erector spinae; cES, Contralateral erector spinae; GMAX, Gluteus maximus; GMED, Gluteus medius; MH, Medial Hamstrings; BF, Biceps Femoris; VM, Vastus Medialis; RF, Rectus Femoris; VL, Vastus Lateralis; AT, Anterior Tibialis; MG, Medial Gastrocnemius.

**Bold text indicates statistically significant difference between the stable and unstable conditions.**

GMED ( $p < 0.001$ ), GMAX ( $p = 0.04$ ), MH ( $p = 0.001$ ), VM ( $p < 0.001$ ), RF ( $p < 0.001$ ), VL ( $p < 0.001$ ), and MG ( $p = 0.004$ ), in the PL direction: iEOB ( $p = 0.02$ ), iES ( $p = 0.003$ ), cES ( $p = 0.015$ ), GMED ( $p = 0.01$ ), BF ( $p = 0.019$ ), VM ( $p = 0.001$ ), RF ( $p = 0.001$ ), VL ( $p < 0.001$ ), and MG ( $p = 0.001$ ), and in the PM direction iEOB ( $p = 0.03$ ), iES ( $p = 0.02$ ), GMED ( $p = 0.001$ ), BF ( $p = 0.007$ ), VM ( $p = 0.006$ ), RF ( $p < 0.001$ ), and VL ( $p = 0.005$ ). No significant differences between the stable and unstable surfaces were observed in iRA (A:  $p = 0.5$ , PL:  $p = 0.06$ , PM:  $p = 0.12$ ), cRA (A:  $p = 0.88$ , PL:  $p = 0.13$ , PM:  $p = 0.053$ ), cEOB (A:  $p = 0.052$ , PL:  $p = 0.11$ , PM:  $p = 0.09$ ), and AT (A:  $p = 0.47$ , PL:  $p = 0.14$ , PM:  $p = 0.7$ ) for any of the directions of the YBT (Table 2). The reach distance between the two conditions for all the three directions were higher for the stable versus the unstable condition (Table 3).

**Table 3. Normalized reach distance during the YBT.**

Condition →	Stable Mean ± SD (% leg length)	Unstable Mean ± SD (% leg length)	% Difference
Directions ↓			
A	63±7	60±7	4.87
PL	82±10	78±9	5.0
PM	89±8	87±9	2.3

SD, Standard Deviation; A, Anterior; PL, Posterolateral; PM, Posteromedial

## DISCUSSION

The present study compared the muscle activation of the various trunk (iRA, cRA, iEOB, cEOB, iES, cES) and LE muscles (GMED, GMAX, MH, BF, VM, RF, VL, AT, MG) while per-

forming the YBT on stable and unstable surfaces. Significant differences in muscle activation were noted for eleven out of fifteen muscles due to the change in the surface. The findings of the current study may provide evidence regarding utilizing the YBT as a training tool in stable and unstable conditions. To the authors' knowledge, no previous studies have measured muscle activation during this test. In general, the addition of unstable surfaces while performing lower extremities weight-bearing exercises have been documented to either increase, decrease or cause no change in the muscle activation patterns.<sup>14-16,18,30-35</sup>

Adding an unstable surface during the performance of the YBT resulted in significantly increased muscle activation in nine (iEOB, iES, cES, GMED, GMAX, BF, VM, RF, VL, MG) out of fifteen muscles in the current study. This is similar to the results found by other authors<sup>18,31</sup> who investigated the effect of various surfaces while performing bridging exercises. To overcome the effects of instability, the body tends to attempt to offer stability by increasing the muscle activation of certain muscles. Alfuth et al.<sup>30</sup> and Ridder et al.<sup>36</sup> reported that adding unstable surfaces while having subjects perform a single leg stance activity caused an increase in the activation of the selected LE muscles. Increased level of external perturbations while maintaining balance on the unstable surfaces requiring more co-contraction of the muscles has been hypothesized as one of the possible reasons for increased stabilizing muscle activation.<sup>32,34</sup> Krause et al.<sup>33</sup> also reported the variations in the selected LE muscle recruitment in the subjects performing a standard lunge compared to the lunge performed in suspension equipment. The suspension lunge condition resulted in increased muscle activation requiring more stability due to the unstable environment created by the suspension equipment. Unstable surfaces have also been reported to cause increased muscle recruitment in the selected LE muscles while performing upper extremity exercises. Shin et al.<sup>16</sup> found that when subjects were asked to perform upper extremity exercises while standing on an unstable surface, there was an increase in the activation of the selected muscles near the ankle joint in an attempt to maintain balance on the unstable surface using secondary to use of an ankle strategy by the subjects.

A large effect size was observed for RF in all the directions, for GMED, VM, and VL in two out of the three directions, and for MG, one out of two directions. A moderate effect size was observed for these muscles in the other directions. The moderate effect size was also observed for the iEOB, iES, and BF for two out of the three directions and cES, GMAX, and MH for one out of the three directions. Moderate to large effect size shows that the higher muscle activity during the unstable surface is due to the difference in condition, and is likely clinically meaningful. In most directions, a large effect size was observed for the quadriceps and GMED muscles because they are key to maintaining balance during a single-leg squat task.<sup>37</sup>

Compared to a stable surface, performing the YBT on an unstable surface resulted in no significant difference in five muscles (iRA, cRA, cEOB, VL, AT) and a decrease in muscle activity for the MH muscle. These findings are also in agreement with the conclusions from previous researchers investigating the effects of various surfaces on muscle recruit-

ment.<sup>15,35</sup> Some of the proposed reasons for no change in muscle activation under unstable conditions included variability in the subjects' training/activity levels, the type of the unstable surfaces used, the distance of the muscle from the unstable surface, the muscle role (prime movers versus stabilizers) in performing the exercise, and biomechanical movement compensations due to added demand on the neuromuscular system.<sup>15,35,38,39</sup>

The use of unstable surfaces has been proposed to place more demands on the neuromuscular system, which in turn may help to improve postural stability and endurance through neural adaptations.<sup>15,40-44</sup> Basketball players, when trained on unstable surfaces, have shown improvement in balance and stability in contrast to their controls who trained on a stable surface.<sup>44</sup> Performing stabilization exercises on unstable surfaces have been reported to be more effective than stable surfaces at improving pain, stability, and disability measured using visual analog scale, Stork Balance Stand test, and Oswestry Disability Index, respectively, in patients with lumbar pain.<sup>41</sup> Performing exercises on unstable surfaces demonstrated greater therapeutic effects in comparison to stable surfaces to improve balance in healthy individuals. The authors suggested that it may be because the unstable surfaces are more effective in influencing the somatosensory system.<sup>45</sup> Exercising on unstable surfaces has been shown to be more effective in improving balance and walking abilities among stroke patients than stable surfaces.<sup>40,42</sup> Authors have also shown the utility of introducing unstable load during strengthening exercises to stress the neuromuscular system.<sup>45-47</sup> The results of their studies showed a decrease in muscle activity for the prime movers but an increase in muscle activity for the stabilizer muscles. They hypothesized that an unstable load places extra demand on the body for stabilization, challenging balance and neuromuscular control.

Ganesh et al.<sup>12</sup> and Chaiwanichsiri et al.<sup>13</sup> utilized the SEBT as an intervention to improve balance and proprioception in their study subjects. Although the YBT is commonly used to measure balance and NM control, based upon the current results, having individuals perform the YBT on the unstable surface may increase muscle activity in key stabilizing muscles and may be considered for use as a training tool for improving balance and stability. However, if clinicians choose to use YBT to improve balance and NM control in their clients, another test and measure should be utilized for pre-post assessment to avoid a threat to construct validity of the YBT via a learning effect.

#### LIMITATIONS AND FUTURE SUGGESTIONS

The use of surface EMG to describe muscular activity during a dynamic activity has limitations such as skin displacement, movement artifact, motor unit recruitment variations, a sub-maximal effort by participants, and the possibility of cross-talk from surrounding muscles in spite of using methodology that is commonly cited in previous research.<sup>17,19,25,27</sup> This study was conducted using healthy adults; therefore, generalizability to a population that is not similar to the subjects in the current study should be avoided. Confirmation of the findings from the present study is necessary for larger and more diverse populations.

Future research should include three-dimensional analysis to learn more about the relationship between joint kinematics and neuromuscular parameters. It would be interesting to investigate if any gender differences exist while performing the YBT. It would also be interesting to see the effect of various types of unstable surfaces on YBT performance.

## CONCLUSION

The results of this study indicate that the performance of the YBT on different surfaces produced a change in the activation of many of the trunk and LE muscles, with the unstable surface inducing increased activation for most of

the studied muscles. Clinically, the results from the current study may guide the use of the YBT with an unstable surface as an exercise intervention for training balance and NM control.

.....

## CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare.

Submitted: April 12, 2021 CDT, Accepted: January 09, 2022 CDT



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc/4.0> and legal code at <https://creativecommons.org/licenses/by-nc/4.0/legalcode> for more information.

## REFERENCES

1. Ruffe NJ, Sorce SR, Rosenthal MD, Rauh MJ. Lower quarter-and upper quarter Y balance tests as predictors of running-related injuries in high school cross-country runners. *Int J Sports Phys Ther.* 2019;14(5):695.
2. Kramer TA, Sacko RS, Pfeifer CE, Gatens DR, Goins JM, Stodden DF. The association between the functional movement screentm, y-balance test, and physical performance tests in male and female high school athletes. *Int J Sports Phys Ther.* 2019;14(6):911.
3. Plisky PJ, Gorman PP, Butler RJ, Kiesel KB, Underwood FB, Elkins B. The reliability of an instrumented device for measuring components of the star excursion balance test. *N Am J Sports Phys Ther.* 2009;4:92-99.
4. Coughlan GF, Fullam K, Delahunt E, Gissane C, Caulfield BM. A comparison between performance on selected directions of the star excursion balance test and the Y balance test. *J Athl Train.* 2012;47(4):366-371.
5. Butler RJ, Lehr ME, Fink ML, Kiesel KB, Plisky PJ. Dynamic balance performance and noncontact lower extremity injury in college football players: an initial study. *Sports Health.* 2013;5(5):417-422.
6. Munro AG, Herrington LC. Between-session reliability of the star excursion balance test. *Phys Ther Sport.* 2010;11(4):128-132.
7. Smith CA, Chimera NJ, Warren M. Association of y balance test reach asymmetry and injury in division I athletes. *Med Sci Sports Exerc.* 2015;47(1):136-141.
8. Butler RJ, Bullock G, Arnold T, Plisky P, Queen R. Competition-level differences on the lower quarter Y-balance test in baseball players. *J Athl Train.* 2016;51(12):997-1002.
9. Hartley EM, Hoch MC, Boling MC. Y-balance test performance and BMI are associated with ankle sprain injury in collegiate male athletes. *J Sci Med Sport.* 2018;21(7):676-680.
10. Gribble PA, Hertel J, Plisky P. Using the Star Excursion Balance Test to assess dynamic postural-control deficits and outcomes in lower extremity injury: a literature and systematic review. *J Athl Train.* 2012;47(3):339-357.
11. Dutton M. *Dutton's Orthopaedic: Examination, Evaluation and Intervention.* 5th ed. McGraw-Hill Education; 2019.
12. Ganesh GS, Chhabra D, Pattnaik M, Mohanty P, Patel R, Mrityunjay K. Effect of trunk muscles training using a star excursion balance test grid on strength, endurance and disability in persons with chronic low back pain. *J Back Musculoskelet Rehabil.* 2015;28(3):521-530.
13. Chaiwanichsiri D, Lorprayoon E, Noomanoch L. Star excursion balance training: effects on ankle functional stability after ankle sprain. *J Med Assoc Thai.* 2005;88:S90.
14. Behm DG, Colado Sanchez JC. Instability resistance training across the exercise continuum. *Sports Health.* 2013;5(6):500-503.
15. Andersen LL, Andersen JL, Magnusson SP, Aagaard P. Neuromuscular adaptations to detraining following resistance training in previously untrained subjects. *Eur J Appl Physiol.* 2005;93(5-6):511-518.
16. Shin D, Cha J, Song C. Electromyographic analysis of trunk and lower extremity muscle activities during pulley-based shoulder exercises performed on stable and unstable surfaces. *J Phys Ther Sci.* 2015;27(1):71-74.
17. Vezina MJ, Hubley-Kozey CL. Muscle activation in therapeutic exercises to improve trunk stability. *Arch Phys Med Rehabil.* 2000;81(10):1370-1379.
18. Czaprowski D, Afeltowicz A, Gebicka A, et al. Abdominal muscle EMG-activity during bridge exercises on stable and unstable surfaces. *Phys Ther Sport.* 2014;15(3):162-168.
19. Norris B, Trudelle-Jackson E. Hip- and thigh-muscle activation during the star excursion balance test. *J Sport Rehabil.* 2011;20(4):428-441.
20. Andersen LL, Magnusson SP, Nielsen M, Haleem J, Poulsen K, Aagaard P. Neuromuscular activation in conventional therapeutic exercises and heavy resistance exercises: Implications for rehabilitation. *Phys Ther.* 2006;86(5):683-697.
21. Bhanot K, Kaur N, Brody LT, Bridges J, Berry DC, Ode JJ. Hip and trunk muscle activity during the Star Excursion Balance Test in healthy adults. *J Sport Rehabil.* 2019;28(7):682-691.
22. Earl JE, Hertel J. Lower-extremity muscle activation during the star excursion balance tests. *J Sport Rehabil.* 2001;10(2):93-104.

23. Thompson WR, Gordon NF, Pescatello LS. *ACSM's Guidelines for Exercise Testing and Prescription*. 8th ed. Lippincott Williams & Wilkins; 2009.
24. Escamilla RF, Babb E, DeWitt R, et al. Electromyographic analysis of traditional and nontraditional abdominal exercises: implications for rehabilitation and training. *Phys Ther*. 2006;86(5):656-671.
25. Cram JR, Criswell E. *Cram's Introduction to Surface Electromyography*. 2nd ed. Jones & Bartlett Publishers; 2010.
26. Fradkin AJ, Gabbe BJ, Cameron PA. Does warming up prevent injury in sport? the evidence from randomised controlled trials. *J Sci Med Sport*. 2006;9(3):214-220. doi:10.1016/j.jsams.2006.03.026
27. Kendall FP, McCreary EK, Provance PG, Rodgers MM, Romani WA. *Muscles: Testing and Function, with Posture and Pain*. 5th ed. Lippincott Williams & Wilkins; 2005.
28. Gribble PA, Hertel J. Considerations for normalizing measures of the Star Excursion Balance Test. *Meas Phys Educ Exerc Sci*. 2003;7(2):89-100.
29. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. Academic Press; 2013.
30. Alfuth M, Gomoll M. Electromyographic analysis of balance exercises in single-leg stance using different instability modalities of the forefoot and rearfoot. *Phys Ther Sport*. 2018;31:75-82.
31. Choi K, Bak J, Cho M, Chung Y. The effects of performing a one-legged bridge with hip abduction and use of a sling on trunk and lower extremity muscle activation in healthy adults. *J Phys Ther Sci*. 2016;28(9):2625-2628.
32. Ferreira LAB, Pereira WM, Rossi LP, Kerpers II, de Paula Jr AR, Oliveira CS. Analysis of electromyographic activity of ankle muscles on stable and unstable surfaces with eyes open and closed. *J Bodyw Mov Ther*. 2011;15(4):496-501.
33. Krause DA, Elliott JJ, Fraboni DF, McWilliams TJ, Rebhan RL, Hollman JH. Electromyography of the hip and thigh muscles during two variations of the lunge exercise: a cross-sectional study. *Int J Sports Phys Ther*. 2018;13(2):137.
34. Lee S, Choi YH, Kim J. The effects of changes in support and inclined boards on lower-extremity muscle activity. *J Phys Ther Sci*. 2017;29(6):1045-1047.
35. Wahl MJ, Behm DG. Not all instability training devices enhance muscle activation in highly resistance-trained individuals. *J Strength Cond Res*. 2008;22(4):1360-1370.
36. De Ridder R, Willems T, Vanrenterghem J, Roosen P. Influence of balance surface on ankle stabilizing muscle activity in subjects with chronic ankle instability. *J Rehabil Med*. 2015;47(7):632-638.
37. Claiborne TL, Armstrong CW, Gandhi V, Pincivero DM. Relationship between hip and knee strength and knee valgus during a single leg squat. *Journal of applied biomechanics*. 2006;22(1):41-50.
38. De Mey K, Danneels L, Cagnie B, et al. Shoulder muscle activation levels during four closed kinetic chain exercises with and without Redcord slings. *J Strength Cond Res*. 2014;28(6):1626-1635.
39. Lehman GJ, Gilas D, Patel U. An unstable support surface does not increase scapulothoracic stabilizing muscle activity during push up and push up plus exercises. *Man Ther*. 2008;13(6):500-506.
40. Bang DH, Shin WS, Noh HJ, Song MS. Effect of unstable surface training on walking ability in stroke patients. *J Phys Ther Sci*. 2014;26(11):1689-1691.
41. Kang TW, Lee J hyun, Park DH, Cynn HS. Effect of 6-week lumbar stabilization exercise performed on stable versus unstable surfaces in automobile assembly workers with mechanical chronic low back pain. *Work*. 2018;60(3):445-454.
42. Kim BS. Effects of pressure sense perception training on unstable surface on somatosensory, balance and gait function in patients with stroke. *J Korean Soc Phys Med*. 2015;10(3):237-245.
43. Nam HC, Cha HG, Kim MK. The effects of exercising on an unstable surface on the gait and balance ability of normal adults. *J Phys Ther Sci*. 2016;28(7):2102-2104.
44. Sedaghati P. The effect of selective plyometric exercises using an unstable surface on the movement performance of basketball players. *Ann Appl Sport Sci*. 2018;6(3):15-22.
45. Kohler JM, Flanagan SP, Whiting WC. Muscle activation patterns while lifting stable and unstable loads on stable and unstable surfaces. *J Strength Cond Res*. 2010;24(2):313-321.
46. Saeterbakken AH, Solstad TEJ, Stien N, Shaw MP, Pedersen H, Andersen V. Muscle activation with swinging loads in bench press. *PLOS One*. 2020;15(9):e0239202.

47. Solstad TE, Andersen V, Shaw M, Hoel EM, Vonheim A, Saeterbakken AH. A Comparison of Muscle Activation between Barbell Bench Press and Dumbbell Flyes in Resistance-Trained Males. *J Sports Sci Med.* 2020;19(4):645.